

“Vermindert pijn”

Romanelli M, Evaluation of the efficacy and tolerability of a solution containing propyl betaine and polyhexanide, *Skin Pharmacol Physiol* 2010;23(Suppl 1):41-44.

Durante C., Evaluation of the effectiveness of a polyhexanide and propylbetaine-based gel in the treatment of chronic wound, *Minerva Chir.*, 2014, 69(5):283-92.

Valenzuela AR., The effectiveness of a 0.1 % polyhexanide gel. *Rev ROL Enf*, 2008;31(4):247-252.

To E., The effectiveness of topical Polyhexamethylene Biguanide (PHMB) Agents for the Treatment of Chronic Wounds: A systematic Review, *Surgical Technology International XXIX* 2016.

Forma, Evaluation of Prontosan wound irrigation solution as a coadjuvant in the treatment of ulcers, *ActaVulnologica* 2016.

Horrocks A., Prontosan® Wound Irrigation Solution and Gel: Management of chronic wounds, *Br J Nurs* 2006;15(22):1222-28.

Cutting K, Addressing the challenge of wound cleansing in the modern era, *British Journal of Nursing*, 2010 (Tissue Viability Supplement), Vol 19, No 11.

Smith S., Successful management of infected wounds using a solution and gel containing Betaine and PHMB, *Wound Practice and Research*, 2013.

Andriessen A., Assessment of a wound cleansing solution in the treatment of problem wounds , *WOUNDS* ;2008, 20(6):171-175.

Cutting K.,Addressing the challenge of wound cleansing in the modern era, *British Journal of Nursing*, (Tissue Viability Supplement), 2010, Vol 19, No 11.

Valenzuela AR., The effectiveness of a 0.1 % polyhexanide gel. *Rev ROL Enf*, 2008;31(4):247-252.

Bellingeri, A., Effect Of A Wound Cleansing Solution On Wound Bed Preparation And Inflammation In Chronic Wounds: A Single-Blind RCT". *Journal of Wound Care*, 2016, 160-168. Web.

“Mag worden toegepast bij pasgeborenen en kinderen”

Dekra Letter of Notification
Poster Presentation Ciprandi

B. Braun Medical B.V. | www.bbraun.com

De informatie in deze communicatie is strikt vertrouwelijk, kan bedrijfsinformatie omvatten en is alleen bedoeld voor gebruik door de geadresseerde. Het is eigendom van de afzender van deze informatie. Ongeoorloofd gebruik, openbaarmaking of kopiëren van deze communicatie of een deel daarvan is ten strengste verboden en kan onwettig zijn.

Claimlist

Neem
wondbedvoorbereiding
serieus

Gebruik Prontosan®



Prontosan®

Voor je ligt de officiële lijst met claims van Prontosan®. Een handig overzicht van alle claims met een verwijzing naar de betreffende studie of wetenschappelijke onderbouwing.



"Voorkomt en behandelt bioflim"

Smith S., Successful management of infected wounds using a solution and gel containing Betaine and PHMB, Wound Practice and Research, 2013.

"Verkort wondgenezingstijd"

Queiros R., The effectiveness of cleansing solutions for wound treatment: a systematic review, JBLROLL, 2013, 11:5 (169-181).

Kiefer J., An observational study with Prontosan® Wound Gel X in partial and full thickness skin burns, Poster Presentation WUWHS, 2016.

Harley J., Evaluation of efficacy and cost saving when including PHMB and Betaine Hydrogel in the treatment of paediatric burns, Poster Presentation, 2015.

Eberlein T., Expert assessment on the benefits of systematic application of Prontosan® solution in wound treatment with particular focus on cost-efficiency when compared to current standard treatment (saline/Ringer), Internal data, 2006.

Andriessen A., Assessment of a wound cleansing solution in the treatment of problem wounds, WOUNDS ;2008, 20(6):171-175.

Cutting K., Addressing the challenge of wound cleansing in the modern era, British Journal of Nursing, (Tissue Viability Supplement), 2010, Vol 19, No 11.

Valenzuela AR., The effectiveness of a 0.1 % polyhexanide gel. Rev ROL Enf, 2008;31(4):247-252.

Bellingeri, A., Effect Of A Wound Cleansing Solution On Wound Bed Preparation And Inflammation In Chronic Wounds: A Single-Blind RCT". Journal of Wound Care, 2016, 160-168. Web.

"Versnelt het wondgenezingsproces"

Bellingeri, A., Effect Of A Wound Cleansing Solution On Wound Bed Preparation And Inflammation In Chronic Wounds: A Single-Blind RCT". Journal of Wound Care, 2016, 160-168. Web.

Roth C., Response analysis of stimulating efficacy of polyhexanide in an in vitro wound model with respiratory ciliary epithelial cells, Skin Pharmacol Physiol, 2010, 23(suppl1):35-40.

To, The effectiveness of PHMB agents for the treatment of chronic wounds a systematic review, Advance wound healing, 2016.

Eberlein T, Clinical use of polyhexanide on acute and chronic wounds for antiseptics and decontamination, Skin Pharmacol Physiol, 2010; 23(suppl1):45-51.

Durante C., Evaluation of the effectiveness of a polyhexanide and propylbetaine-based gel in the treatment of chronic wound, Minerva Chir., 2014, 69(5):283-92.

Moeller A, Experiences in using polyhexanide containing wound products in the management of chronic wounds – results of a methodical and retrospective analysis of 953 cases. Wundmanagement; 2008,3:112-117.

Horrocks A., Prontosan® Wound Irrigation Solution and Gel: Management of chronic wounds, Br J Nurs 2006;15(22):1222-28.

"Verbeter de kwaliteit van leven"

Harley J., Evaluation of efficacy and cost saving when including PHMB and Betaine Hydrogel in the treatment of paediatric burns, Poster Presentation, 2015

Valenzuela AR., The effectiveness of a 0.1 % polyhexanide gel. Rev ROL Enf, 2008;31(4):247-252.

Posnett, The resource impact of wounds on health-care providers in Europe, Journal of wound care, 2009

Cutting K., Addressing the challenge of wound cleansing in the modern era, British Journal of Nursing, (Tissue Viability Supplement), 2010, Vol 19, No 11.

Horrocks A., Prontosan® Wound Irrigation Solution and Gel: Management of chronic wounds, Br J Nurs 2006;15(22):1222-28.

"Vermindert geur"

Romanelli M., Evaluation of the efficacy and tolerability of a solution containing propyl betaine and polyhexanide, Skin Pharmacol Physiol 2010;23(Suppl 1):41-44.

Moeller A, Experiences in using polyhexanide containing wound products in the management of chronic wounds – results of a methodical and retrospective analysis of 953 cases. Wundmanagement; 2008,3:112-117.

Valenzuela AR., The effectiveness of a 0.1 % polyhexanide gel. Rev ROL Enf, 2008;31(4):247-252.

Forma, 2006, Evaluation of Prontosan wound irrigation solution as a coadjuvant in the treatment of ulcers, ActaVulnologica

Horrocks A., Prontosan® Wound Irrigation Solution and Gel: Management of chronic wounds, Br J Nurs 2006;15(22):1222-28.

Queiros R., The effectiveness of cleansing solutions for wound treatment: a systematic review, JBLROLL, 2013, 11:5 (169-181).

"Effectiever dan NaCl 0,9%"

Bellingeri, A., Effect Of A Wound Cleansing Solution On Wound Bed Preparation And Inflammation In Chronic Wounds: A Single-Blind RCT". Journal of Wound Care, 2016, 160-168.

Andriessen A., Assessment of a wound cleansing solution in the treatment of problem wounds, WOUNDS ;2008, 20(6):171-175.

Romanelli M, Evaluation of the efficacy and tolerability of a solution containing propyl betaine and polyhexanide, Skin Pharmacol Physiol 2010;23(Suppl 1):41-44.

Davis SC, "Effectiveness of a polyhexanide irrigation solution on methicillin-resistant Staphylococcus aureus biofilms in a porcine wound model" in International Wound Journal ISSN 1742-4801, 2017, 1-8, © 2017 Medicalhelplines.com Inc and John Wiley & Sons Ltd.

Cutting K, Addressing the challenge of wound cleansing in the modern era, British Journal of Nursing, 2010 (Tissue Viability Supplement), Vol 19, No 11.

Seipp HM, Efficacy of various wound irrigation solutions against biofilms, Zeitschrift für Wundheilung (ZFW), 2005; 4: 160-164.